AMENDMENTS TO THE CLAIMS

1-6. (Cancelled)

7. (Previously presented) A percutaneous absorption preparation comprising (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno [5,4-b]furan-8-yl)ethyl]acetamide, lauric diethanolamide, and optionally one or more members selected from fatty acid esters and polyhydric alcohols.

8-19. (Cancelled)

- **20.** (**Previously presented**) A percutaneous absorption preparation comprising (S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno[5,4-b]furan-8-yl)ethyl]acetamide, isopropyl myristate, polyethylene glycol and lauric diethanolamide.
- 21. (Currently amended) A percutaneous absorption preparation comprising a compound having a melatonin receptor agonist activity, lauric diethanolamide and optionally one or more members selected from fatty acid esters and polyhydric alcohols, wherein the compound having a melatonin receptor agonist activity is a compound represented by the formula:

$$\begin{array}{c|c}
R^{2} \\
N \\
N \\
R^{1}
\end{array}$$

$$\begin{array}{c|c}
R^{1} \\
R^{2} \\
N \\
R^{3}
\end{array}$$

wherein, R¹ represents an optionally substituted hydrocarbon group, an optionally substituted amino group or an optionally substituted heterocyclic group;

R² represents a hydrogen atom or an optionally substituted hydrocarbon group;

R³ represents a hydrogen atom, an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group;

X represents CHR⁴, NR⁴[[,]] or O or S in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group;

Y represents C[[,]] or CH or N, provided that when X is CH₂, Y is C or CH;

represents a single bond or a double bond;

ring A represents an optionally substituted, 5- to 7-membered oxygen-containing heterocyclic ring;

ring B represents an optionally substituted benzene ring; and

m represents an integer of 1 to 4;

or a salt thereof, wherein the percutaneous absorption preparation is a skin plaster or a skin patch which is applied and/or attached to the skin.

22-32. (Cancelled)

33. (Currently amended) A percutaneous absorption preparation comprising a compound having a melatonin receptor agonist activity, lauric diethanolamide and optionally one or more members selected from fatty acid esters and polyhydric alcohols, wherein the compound having a melatonin receptor agonist activity is a compound represented by the formula:

$$\begin{array}{c|c}
R^{2} \\
N \\
N \\
R^{1}
\end{array}$$

$$\begin{array}{c|c}
R^{1} \\
R^{2} \\
N \\
R^{3}
\end{array}$$

wherein, R¹ represents an optionally substituted hydrocarbon group, an optionally substituted amino group or an optionally substituted heterocyclic group;

R² represents a hydrogen atom or an optionally substituted hydrocarbon group;

R³ represents a hydrogen atom, an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group;

X represents CHR⁴, NR⁴[[,]] or O or S in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group;

Y represents C[[,]] or CH or N, provided that when X is CH₂, Y is C or CH;

represents a single bond or a double bond:

ring A represents an optionally substituted, 5- to 7-membered oxygen-containing heterocyclic ring;

ring B represents an optionally substituted benzene ring; and

m represents an integer of 1 to 4;

or a salt thereof, wherein the percutaneous absorption preparation is contained in a skin contact member comprising silicon dioxide.

34-38. (Cancelled)

39. (Currently amended) A method of treating diseases related to melatonin, which comprises administrating administering to a patient with a melatonin related disease a percutaneous absorption preparation comprising a compound having a melatonin receptor agonist activity, lauric diethanolamide and optionally one or more members selected from fatty acid esters and polyhydric alcohols, wherein the compound having a melatonin receptor agonist activity is a compound represented by the formula:

$$\begin{array}{c|c}
R^{2} \\
N \\
R^{1}
\end{array}$$

$$\begin{array}{c|c}
R^{1} \\
R^{2} \\
R^{3}
\end{array}$$

wherein, R¹ represents an optionally substituted hydrocarbon group, an optionally substituted amino group or an optionally substituted heterocyclic group;

R² represents a hydrogen atom or an optionally substituted hydrocarbon group;

R³ represents a hydrogen atom, an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group;

X represents CHR⁴, NR⁴[[,]] or O or S in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group;

Y represents C[[,]] or CH or N, provided that when X is CH2, Y is C or CH;

represents a single bond or a double bond;

ring A represents an optionally substituted, 5- to 7-membered oxygen-containing heterocyclic ring;

ring B represents an optionally substituted benzene ring; and m represents an integer of 1 to 4; or a salt thereof.

40. (Currently amended) A method for percutaneous absorption of a compound having a melatonin receptor agonist activity, which comprises administering to a patient with a melatonin related disease a percutaneous absorption preparation comprising a compound having a melatonin receptor agonist activity, lauric diethanolamide and optionally one or more members selected from fatty acid esters and polyhydric alcohols, wherein the compound having a melatonin receptor agonist activity is a compound represented by the formula:

$$\begin{array}{c|c}
R^{2} \\
N \\
N \\
R^{1}
\end{array}$$

$$\begin{array}{c|c}
R^{1} \\
R^{2} \\
N \\
R^{3}
\end{array}$$

wherein, R¹ represents an optionally substituted hydrocarbon group, an optionally substituted amino group or an optionally substituted heterocyclic group;

R² represents a hydrogen atom or an optionally substituted hydrocarbon group;

R³ represents a hydrogen atom, an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group;

X represents CHR⁴, NR⁴[[,]] or O or S in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group;

Y represents C[[,]] or CH or N, provided that when X is CH₂, Y is C or CH;

represents a single bond or a double bond;

ring A represents an optionally substituted, 5- to 7-membered oxygen-containing heterocyclic ring;

ring B represents an optionally substituted benzene ring; and m represents an integer of 1 to 4; or a salt thereof.

41. (Cancelled)

- **42.** (**Previously presented**) The method according to claim 39, wherein the percutaneous absorption preparation is affixed between about 6 hours before bedtime to just before bedtime.
- **43.** (New) The percutaneous absorption preparation according to claim 21, wherein X represents CHR⁴ in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group.

- **44.** (New) The percutaneous absorption preparation according to claim 33, wherein X represents CHR⁴ in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group.
- 45. (New) The method of treating diseases related to melatonin according to claim 39, wherein X represents CHR⁴ in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group.
- **46.** (New) The method for percutaneous absorption of a compound according to claim 40, wherein X represents CHR⁴ in which R⁴ represents a hydrogen atom or an optionally substituted hydrocarbon group.